

Year	Age	Sex	Height	Weight	Body Mass Index	Waist Circumference	Waist-Hip Ratio	Trunk Fat (%)	Visceral Fat (cm ²)	Subcutaneous Fat (cm ²)	Visceral Fat Index	Subcutaneous Fat Index	Visceral Fat to Subcutaneous Fat Ratio
1990	20	Male	170	70	24.0	85	0.85	15	10	5	0.10	0.05	2.0
1990	20	Female	160	55	21.5	75	0.80	12	8	4	0.08	0.04	2.0
1995	25	Male	175	80	25.9	90	0.88	18	12	6	0.12	0.06	2.0
1995	25	Female	165	65	23.8	80	0.82	14	10	5	0.10	0.05	2.0
2000	30	Male	180	90	27.8	95	0.90	20	15	7	0.15	0.07	2.1
2000	30	Female	170	75	25.9	85	0.85	16	12	6	0.12	0.06	2.0
2005	35	Male	185	100	29.4	100	0.92	22	18	8	0.18	0.08	2.2
2005	35	Female	175	85	27.9	90	0.88	18	14	7	0.14	0.07	2.0
2010	40	Male	190	110	31.0	105	0.94	24	20	9	0.20	0.09	2.2
2010	40	Female	180	95	29.4	95	0.90	20	16	8	0.16	0.08	2.0
2015	45	Male	195	120	31.2	110	0.96	26	22	10	0.22	0.10	2.2
2015	45	Female	185	105	29.9	100	0.92	22	18	9	0.18	0.09	2.0
2020	50	Male	200	130	32.5	115	0.98	28	24	11	0.24	0.11	2.2
2020	50	Female	190	115	31.6	105	0.94	24	20	10	0.20	0.10	2.0

4

Priority Claimed:: Yes